

## ISOLATION, PHYSICO-CHEMICAL AND BIOLOGICAL PROPERTIES OF THE IMMUNITY POLYPEPTIDE BIOREGULATOR FROM THYMUS

O.A.Pisarev, V.G.Morosov, V.Kh.Khavinson, L.K.Shataeva, G.V. Samsonov

Institute of Macromolecular Compounds, Academy of Sciences of the USSR, Leningrad

### Introduction

The main subject of modern immunology is to investigate the ways and means for the recovery of the disturbed immunological functions in the process of human vital activity. One of the hypotheses accounting for a T-lymphocyte formation is the assumption that in thymus a hormonal factor is produced, due to its action the marrow stem cells obtain T-lymphocyte properties and are further differentiated into T-cells participating in immunological reactions (1). Thymosin is a well-known isolated thymus factor (2). There are some reports on the thymic humoral factor isolation (3), thymopoietin (4) as well as the circulating blood thymus factor (5). This study had the aim of the isolation of a polypeptide fraction stimulating immunogenesis and the examination of its composition, physico-chemical and biological properties.

### Results

The main stages of the compound production from thymus are represented in Fig.1. Before the stage of the ion-exchange chromatography more than 10 compounds of different origin were present. The multi-component extract was fractionated on Bio-carb (6-8) into three fractions; electrophoresis on polyacrylamide gel showed that in fraction one there are nine com-

pounds, in fraction 2 3 compounds and in fraction 3 there is 1 compound present. Fraction 1 eluates from the ion-exchange column at pH 3.0, fraction 2 at pH 6.5, fraction 3 at pH 9.5 - 10.0.

thymus tissue  
 acetone  
 (-4°, 48h)  
 homogenization  
 extraction  
 (6 x 3 % CH<sub>3</sub>COOH + ZnCl<sub>2</sub>, 72 h)  
 centrifugation  
 acetone  
 (-4°, 1 h)  
 dissolution in 5 % CH<sub>3</sub>COOH  
 ion-exchange chromatography (Biocarb-T)  
 thymalin

Fig.1. Scheme of the isolation of thymalin.

According to gel chromatography on Sephadex G-25 and G-10 fraction 3 was also homogeneous. Investigations by ultracentrifugation demonstrated a symmetrical peak for fraction 3 with the sedimentation coefficient 0.7 Sw.

The percentages of each fraction, molecular masses, isoelectric points, tyrosine contents (by Lowry) and also the results of biological testing are shown in Figs. 2 and 3.

Fraction 1 has no biological activity. Fraction 2 stimulated immunological reactions 15-20 times stronger than compared to the extract. The biological activity of the extract is associated mainly with fraction 3. Intramuscular injection (0.5 mg/kg) of this fraction to the mouse increased the number of lymphocytes in blood by 50-70 %, and 0.1-0.5 mg/kg stimulated the formation of antibody forming cells threefold. At higher dose (25 mg/kg) this fraction had no stimulating effect and in some cases suppressed immunological reactions. Fraction 3 was named "thymalin" and was later used for clinical and bio-

logical tests.

Preparation	Dose mg/ kg	Number of lym- phocy- tes	Number of circula- ting an- tibody	Number of an- tibody form- ing cells	Phago- cytic reac- tion	Activity of pero- xidase
Fraction 1	25	-	-	-	-	-
	2.5	-	-	-	-	-
Fraction 2	25	<u>+</u>	<u>+</u>	-	<u>+</u>	<u>+</u>
	2.5	-	-	-	-	-
Fraction 3	25	+	<u>+</u>	-	<u>+</u>	<u>+</u>
	0.5	+	+	+	+	+
	0.1	+	<u>+</u>	+	<u>+</u>	+

Fig. 2. Effect of thymus preparations on the indices of immunologic reactivity.

Component	Content in ex- tract (%)	Mol.	Isoelectr. points	Tyrosine content by Lowry %
Fraction 1	80	700	2.5	4.5
Fraction 2	15	5000	4.5	1.0
Fraction 3	5	5000	9.5	1.0

Fig. 3. Properties of thymic fractions.

Study of the thymalin amino acid composition confirmed its polypeptide structure. Fig.4 shows that thymalin is a basic polypeptide, consisting of 38 amino acid residues. In the thymalin structure there are 11 basic and 4 acid amino acids. N-terminal analysis performed according to (9) demonstrated that N-terminal amino acids are lysine and alanine.

Indices	Thymalin properties
Mol.	50 00
Isoelectric point	9.5
Number of amino acid residues	38
Amino acid composition	Asp <sub>2</sub> , Thr <sub>2</sub> , Ser <sub>2</sub> , Glu <sub>2</sub> , Trp <sub>1</sub> , Pro <sub>3</sub> , Gly <sub>3</sub> , Ala <sub>4</sub> , Val <sub>1</sub> , Met <sub>1</sub> , Ile <sub>1</sub> , Leu <sub>1</sub> , Tyr <sub>1</sub> , Cys <sub>1</sub> , Phe <sub>1</sub> , His <sub>1</sub> , Lys <sub>8</sub> , Arg <sub>3</sub>

Fig. 4. Physico-chemical properties of thymalin.

Conformational properties of thymalin were examined by the ORD method. Thymalin undergoes a cooperative conformational transition in the 8.5 - 10.5 range with  $[\alpha]_D^{410^\circ}$  changing from  $-230^\circ$  to  $-165^\circ$ . Estimates according to (10,11) reveal no ordered structure in the pH area of 4.0-8.5, whereas at pH 10.5 ca. 20 %  $\alpha$ -helix is present.

The results obtained in the study of the thymalin effect on the immunity indices are given in Fig.5. The number of T- and B-lymphocytes was determined by E-POK and EAG-POK content in the lymphatic nodules and the spleen of the animals as described in (12). After thymectomy the number of caryocytes, T-lymphocytes in the spleen, has decreased to a great extent. Cell composition in the lymphatic nodules has not changed. After thymalin injection the number of caryocytes has completely recovered and the number of T- and B-lymphocytes has enhanced in the spleen. In the lymphatic nodules, the changes of caryocyte content, T- and B-lymphocytes appears less noticeable.

The effect of thymalin on T-cell population was confirmed by the investigation of its action in the culture of lymphocytes produced from thymectomized and pseudo-operated guinea pigs. The percentage of T-cells in the spleen months after the operation was  $28.6 \pm 2.9$  for the pseudo-operated animals

Group of animals	Number of caryocytes mln		Number of T-lymphocytes th/mg		Number of B-lymphocytes th/mg	
	Spleen	Lymphatic nodules	Spleen	Lymphatic nodules	Spleen	Lymphatic nodules
Pseudo-operated	58.5 <sup>+</sup> <sub>18.3<sup>-</sup></sub>	84.3 <sup>+</sup> <sub>7.2<sup>-</sup></sub>	158.4 <sup>+</sup> <sub>9.4<sup>-</sup></sub>	310.3 <sup>+</sup> <sub>19.9<sup>-</sup></sub>	342.5 <sup>+</sup> <sub>20.1<sup>-</sup></sub>	362.3 <sup>+</sup> <sub>27.3<sup>-</sup></sub>
Thymectomized	225.1 <sup>+</sup> <sub>16.2<sup>-</sup></sub>	69.1 <sup>+</sup> <sub>4.8<sup>-</sup></sub>	69.2 <sup>+</sup> <sub>12.0<sup>-</sup></sub>	278.1 <sup>+</sup> <sub>25.1<sup>-</sup></sub>	275.4 <sup>+</sup> <sub>23.2<sup>-</sup></sub>	372.2 <sup>+</sup> <sub>31.1<sup>-</sup></sub>
Thymectomized + thymalin	352.6 <sup>+</sup> <sub>16.8<sup>-</sup></sub>	85.2 <sup>+</sup> <sub>5.7<sup>-</sup></sub>	153.2 <sup>+</sup> <sub>7.6<sup>-</sup></sub>	295.5 <sup>+</sup> <sub>21.6<sup>-</sup></sub>	301.3 <sup>+</sup> <sub>25.8<sup>-</sup></sub>	385.2 <sup>+</sup> <sub>30.9<sup>-</sup></sub>

Fig. 5. Effect of the thymus factor on the indices of guinea pig immunity.

and  $16.8 \pm 20$  for the thymectomized ones. When the thymus factor was applied the content of T-cells in the lymphocyte culture of thymectomized guinea pigs increased to the initial indices. The content of T-cells in the lymphocyte culture of the pseudooperated animals did not markedly change.

From the above data we assume that the decrease of the number of T-cells in the spleen after thymectomy occurs owing to  $T_1$ -population of lymphocytes in mature animals which are present mainly in thymus and spleen. Thymectomy of mature animals only slightly affects the cell composition of the lymphatic nodules containing mostly  $T_2$  population of lymphocytes. Thus one can suggest that thymalin recovers the number of T-lymphocytes possessing a function of a helper in a certain stage of their development and thereby favours the definition of antigens and the induction of the cooperative immune response to thymus-dependent antigens. Due to the lack of thymus factor after thymectomy the number of T-lymphocytes decreases, this being the possible reason for suppressed immunological reactivity of thymectomized animals.

Thymus-dependence of lymphocyte population as well as the helper activity of antigene-activated cells are evidently mediated by the presence of thymus factor on the surface of lymphocytes. Hence we may suggest that T-lymphocytes are the carriers of a thymus factor inducing a cooperative immune response to thymus-dependending antigenes. The effect of the thymus factor on different physiological systems is likely to be associated with the function of T-cells which contain genetic information and they are the participants not only of immunologic but also of metabolic processes, such as the regeneration of tissues, the synthesis of protein and nucleic acids, carbohydrate and fat metabolism.

#### References

1. Burnet, F.M.: *Lancet* 1, 358-360 (1970).
2. Goldstein, A.L., Guna, A.: *Proc.Natl.Acad.Sci.* 69, 1800-1810 (1972).
3. Trainin, N., Small, M.: *J.Exp.Med.* 132, 885-897 (1970).
4. Basch, R., Goldstein, G.: *Proc.Natl.Acad.Sci.* 71, 1474-1478 (1974).
5. Bach, A.L., Dardenne, M., Pleau, M.A., Bach, M.A.: *Ann.N.Y.Acad.Sci.* 249, 186-210 (1975).
6. Morozov, V.G., Khavisnon, V.Kh., Pisarev, O.A.: *Proc.Acad.Sci. USSR.* 233, 491-494 (1977).
7. Pisarev, O.A., Shataeva, L.K., Samsonov, G.V.: *Bull.Acad.Sci., USSR.* 29(1), 37-42 (1980).
8. Pisarev, O.A., Shataeva, L.K., Samsonov, G.V.: *J.Physical Chemistry, USSR.* 54. 2391-2395 (1980).
9. Cray, W.R.: *Methods in Enzymol.* 11, 469-475 (1967).
10. Moffit, W., Yang, J.T.: *Proc.Natl.Acad.Sci.* 42, 596-600 (1956).
11. Shechter, E., Blout, E.: *Proc.Natl.Acad.Sci.* 51, 794-798 (1964).
12. Stadecker, M.J., Bishop, G., Wortiz, H.H.: *J.Immunol.* 3, 1834-1837 (1973).