

*Russian Original Vol. 130, No. 11, November, 2000*

---

April, 2001

BULLETIN OF  
**EXPERIMENTAL**  
**BIOLOGY**  
AND **MEDICINE**

**БЮЛЛЕТЕНЬ ЭКСПЕРИМЕНТАЛЬНОЙ  
БИОЛОГИИ И МЕДИЦИНЫ**

**(BYULLETEN' ÉKSPERIMENTAL'NOI  
BIOLOGII I MEDITSINY)**

**TRANSLATED FROM RUSSIAN**

**CONSULTANTS BUREAU, NEW YORK**

## BIOGERONTOLOGY

# Effect of Epithalone on the Age-Specific Changes in the Time Course of Lipid Peroxidation in *Drosophila melanogaster*

V. Kh. Khavinson and S. V. Myl'nikov\*

Translated from *Byulleten' Eksperimental'noi Biologii i Meditsiny*, Vol. 130, No. 11, pp. 585-588, November, 2000  
Original article submitted November 4, 2000

The effect of epithalone on the age-specific time course of lipid peroxidation was studied in inadaptive *Drosophila melanogaster* strains. A single dose of epithalone at the larval stage decreased the level of conjugated hydroperoxides and Schiff bases throughout the life span of imago. Strain- and sex-specific differences in the time course of the studied sign are characterized.

**Key Words:** lipid peroxidation; ageing; epithalone; peptides

Lipid peroxidation (LPO) is the main source of free radicals in tissues. According to the free-radical theory of aging [10], products of these reactions accumulate with age, while inhibition of LPO reactions with antioxidants prolongs the life span. This relationship was demonstrated in many experiments [7-9,12-14].

It was shown that epithalamine, a peptide preparation of the pineal gland, inhibits LPO processes, increases total serum antioxidant activity in rats, and increases serum activity of Cu,Zn-superoxide dismutase and ceruloplasmin concentration [1,3]. Epithalamine inhibits LPO processes in *D. melanogaster* [2].

Previously [10] we characterized the antioxidative properties of epithalamine by measuring the intensity of LPO and activity of antioxidative enzymes in *Drosophila* at 1-2 age points. In this study we analyzed age-specific changes in the levels of lipid peroxides after treatment with epithalone (Ala-Glu-Asp-Glu)

created on the basis of amino acid composition of epithalamine [6]. Epithalone was synthesized at the St. Petersburg Institute of Bioregulation and Gerontology (North-Western Division of Russian Academy of Medical Sciences).

### MATERIALS AND METHODS

Highly inbred strains of *D. melanogaster* selected by low mating capacity of males (HA<sup>-</sup>, BA<sup>-</sup>) [5], characterized by high level of LPO and short life span were used in the study.

Epithalone was added to nutrient medium in a dose of 0.00001% (w/w) at larval stages II-III, so that the exposure did not exceed 2 days. In the control, normal saline was added into the medium. LPO products in flies of different age were extracted from tissue homogenates by heptane:isopropanol mixture (1:1) containing hydroxytoluene antioxidant. The content of primary LPO products (conjugated hydroperoxides, CHP) was estimated by optical density at 233 nm on an SPG-55 spectrophotometer and the content of LPO end-products (Schiff bases, SB) by fluorescence intensity in the blue band spectrum on a Hitachi fluoro-

St. Petersburg Institute of Bioregulation and Gerontology, North-Western Division of Russian Academy of Medical Sciences; \*Department of Genetics and Selection, St. Petersburg State University. Address for correspondence: vvm@medport.ru. Khavinson V. Kh.




Fig. 1. Effect of epithalone on the content of conjugated hydroperoxides in *D. melanogaster* tissue homogenates. Here and in Fig. 2: a, b) BA<sup>-</sup> strain, females and males, respectively; c, d) HA<sup>-</sup> strain, females and males, respectively. Dark circles: control; light circles: experiment; regression lines: 1) control; 2) experiment.

meter [14]. The values were correlated to the fly weight in the sample. Each experimental point is the mean of 3 measurements, 20-30 flies per sample. The data were processed by dispersion and regression analysis [4].

## RESULTS

Epithalone inhibited LPO processes. In the BA<sup>-</sup> strain significant differences between the experimental and control flies were observed starting from the age of 28 days (Table 1). In the HA<sup>-</sup> strain significant differences appeared at different terms depending on the fly gender (Table 1).

Age-specific changes in LPO in all variants of the experiment are best approximated by an equation  $Y=A+B \times \ln t$  (Fig. 1 and 2). The increase of LPO inten-

sity depends on the strain, gender, and nutrition of flies and is determined by a regression coefficient. Dispersion analysis showed that variability of this coefficient for CHP is by 37% determined by strain differences, by 38% by the gender, and by 20% by nutrition of larvae, while for SB, strain differences are responsible for 0.2% total variability, gender for 20%, and nutrition for 40%. Hence, epithalone prevents mainly accumulation of LPO end products.

The dose of epithalone used in this experiment was 100-1000 times lower than the doses prolonging the life span or decreasing the level of free radicals in drosophila [9]. The effect of epithalone was delayed in adult flies in some experimental series. It seems that epithalone is the signal molecule triggering a cascade of regulatory processes, activating the antioxidant sys-




Fig. 2. Effect of epithalone on the content of Schiff bases in *D. melanogaster* tissue homogenates.

TABLE 1. Content of LPO Products in *D. melanogaster* Tissues Treated with Epithalone (% of Control,  $M \pm m$ )

Parameter		Age, days					
		3	6	12	28	34	41
Content of CHP							
BA <sup>-</sup> strain	females	93±5	98±3	96±3	80±2*	85±3*	90±2***
	males	106±8	100±6	97±4	100±6	100±6	94±4
HA <sup>-</sup> strain	females	83±10	70±6**	91±6	94±6	78±4**	72±5*
	males	102±11	78±6**	85±5***	89±7	—	—
Content of SB							
BA <sup>-</sup> strain	females	75±4*	97±4	92±3	81±3*	88±2**	85±2*
	males	98±5	96±4	97±3	88±3**	85±3*	81±2*
HA <sup>-</sup> strain	females	88±5	91±4	88±3***	88±3**	81±2*	72±2*
	males	86±3*	82±2*	90±1*	92±2**	—	—

Note. \* $p < 0.001$ , \*\* $p < 0.01$ , \*\*\* $p < 0.05$  compared to the control.

tem, which manifests by delayed accumulation of LPO products with age.

## REFERENCES

1. V. N. Anisimov, A. V. Arutyunyan, and V. Kh. Khavinson, *Dokl. Ros. Akad. Nauk*, **352**, No. 6, 831-833 (1997).
  2. V. N. Anisimov, S. V. Myl'nikov, T. I. Oparina, and V. Kh. Khavinson, *Ibid.*, **352**, No. 5, 704-707 (1997).
  3. V. N. Anisimov, V. M. Prokopenko, and V. Kh. Khavinson, *Ibid.*, **343**, No. 4, 557-559 (1995).
  4. N. V. Glotov, L. A. Zhivotovskii, M. I. Khovanov, *et al.*, *Biometry* [in Russian], Leningrad (1982).
  5. L. Z. Kaidanov, S. V. Myl'nikov, O. V. Iovleva, *et al.*, *Genetika*, **30**, No. 8, 1085-1096 (1994).
  6. V. G. Morozov and V. Kh. Khavinson, *Uspekhi Sovrem. Biol.*, **96**, No. 6, 339-352 (1983).
  7. Kh. K. Muradyan and V. E. Sabko, *Zh. Evolyuts. Biokhim.*, **25**, No. 4, 431-435 (1989).
  8. S. V. Myl'nikov, A. N. Smimova, and T. I. Oparina, *Genetika*, **30**, No. 11, 1466-1470 (1994).
  9. N. Sh. Nakaidze, V. I. Kuz'min, E. G. Rozantsev, and L. K. Obukhova, *Izv. Akad. Nauk SSSR*, No. 6, 926-929 (1979).
  10. V. Kh. Khavinson and S. V. Myl'nikov, *Byull. Eksp. Biol. Med.*, **129**, No. 4, 420-422 (2000).
  11. D. Harman, *Proc. Natl. Acad. Sci. USA*, **78**, 7124-7128 (1981).
  12. M. L. Heidrick, L. C. Hendricks, and D. E. Cook, *Mech. Ageing Dev.*, **27**, 341-358 (1984).
  13. K. Gonca Akbulut, I. B. Gonu, and H. Akbulut, *Gerontology*, **45**, No. 2, 67-71 (1999).
  14. R. Pamplona, M. Portero-Otin, C. Ruiz, *et al.*, *Mech. Ageing Dev.*, **112**, No. 3, 169-183 (2000).
-