

Epitalon and colon carcinogenesis in rats: Proliferative activity and apoptosis in colon tumors and mucosa

GEORGE KOSSOY¹, JUDIT ZANDBANK², EUGENIE TENDLER³, VLADIMIR ANISIMOV^{4,5},
VLADIMIR KHAVINSON⁵, IRINA POPOVICH⁴, MARK ZABEZHINSKI⁴,
ITSHAK ZUSMAN⁶ and HERZL BEN-HUR¹

¹Laboratory of Experimental Medicine, Rehovot; ²Institute of Pathology, Assaf Harofeh Medical Center, Beer Yaakov-Tserifin; ³Department of Clinical Biochemistry, Rambam Medical Center, Haifa, Israel;

⁴Department of Carcinogenesis and Oncogerontology, NN Petrov Research Institute of Oncology; ⁵St. Petersburg Institute of Bioregulation and Gerontology, St. Petersburg, Russia; ⁶Koret School of Veterinary Medicine,

Faculty of Agricultural, Food and Environmental Quality Sciences, The Hebrew University of Jerusalem, Rehovot, Israel

Received March 24, 2003; Accepted May 21, 2003

Abstract. The effect of the synthetic pineal peptide Epitalon® (Ala-Glu-Asp-Gly) on proliferative activity in colon tumors, and in mucosal epithelial cells adjacent to and located far from tumors was studied in rats. To evaluate the effect of Epitalon on different stages of carcinogenesis, different treatment regimens were used: during the tumor initiation stage, during the tumor-promotion stage, or during the entire process of tumor development. Eighty 2-month-old male LIO rats were exposed weekly to five subcutaneous injections of 1,2-dimethylhydrazine (DMH) at a single dose of 21 mg/kg body weight. Rats were divided into four groups. Control rats (group 1) received saline at a dose of 0.1 ml during the entire experiment. Rats in group 2 were treated with Epitalon at a dose of 1 µg, five times a week, for 6 months, from the first injection of DMH till the end of the experiment. Rats in group 3 were treated with Epitalon after termination of the carcinogen injections. Rats in group 4 were treated with Epitalon only during the period of DMH exposure (for the first 5 weeks of the experiment). DMH induced proliferation of the secretory epithelium, and this phenomenon was accompanied by a decrease in the size of the stromal area and the area of lymph infiltration in colon tumors and in the colon mucosa adjacent to the tumors (group 1). Epitalon attenuated this effect, especially when the treatment was continued throughout the experiment (group 2). It increased the stromal areas, as well as that of lymphoid infiltration in

the colon mucosa adjacent to the tumors. The intensity of lymphoid infiltration was activated in both the colon mucosa adjacent to a tumor and in the tumor. Mitotic activity of tumor cells was significantly inhibited by Epitalon when the treatment was given throughout the experiment (group 2). In parallel, a high level of apoptosis was seen in the same group. Thus, the strongest inhibitory effect of Epitalon on carcinogenesis in the colon mucosa was manifested when the treatment was continued throughout the experiment.

Introduction

The pineal indole hormone melatonin (N-acetyl-5-methoxytryptamine) and the pineal peptide preparation Epithalamin have been shown to inhibit the development of spontaneous and chemically-induced neoplasms (1-3). The synthetic pineal tetrapeptide Epitalon® (Ala-Glu-Asp-Gly), which was recently structured on the basis of the amino acid sequence of Epithalamin, has also been found to reduce the incidence of spontaneous tumors in mice and chemically-induced colon tumors in rats (4,5). However, the mechanism responsible for these changes has only been partially studied.

Experimental colon cancers are manifested by higher proliferative activity and dramatically reduced apoptosis in the colonic mucosa (6). Melatonin has been shown to inhibit cellular proliferation in the rodents' colon (7). Thus, it can be suggested that the inhibitory effect of melatonin on DMH-induced colon carcinogenesis is realized, at least in part, by its influence on tissue homeostasis. Indeed, melatonin has been shown to affect apoptosis and the proliferative properties of normal and tumorous colonic tissues and several splenic lymphoid elements (8,9). The inhibitory effect of Epitalon on bowel carcinogenesis is comparable to that of melatonin (1,10).

In the present study, we compared the effects of DMH and Epitalon on the proliferative activity of tumor cells and mucosal epithelial cells located adjacent to, and far from the tumors, in an attempt to understand the mechanism underlying the oncogenic properties of Epitalon. To evaluate the effect

Correspondence to: Professor I. Zusman, The Koret School of Veterinary Medicine, The Hebrew University of Jerusalem, P.O. Box 12, Rehovot 76100, Israel
E-mail: zusmani@agri.huji.ac.il

Key words: apoptosis, colon cancer, dimethylhydrazine, Epitalon, proliferative activity





Figure 1. Well-differentiated adenocarcinoma of the colon obtained from a control rat exposed to DMH. Note the different phases of mitosis: metaphase (A) and anaphase (B). H&E, x800.

of Epitalon on different stages of carcinogenesis, different regimens of Epitalon treatment were used: during the tumor-initiation stage, during the tumor-promotion stage, or during the entire process of tumor development (5).

Materials and methods

Chemicals. 1,2-Dimethylhydrazine dihydrochloride (DMH) was purchased from Sigma Chemical Co. (St. Louis, MO). The synthetic pineal peptide Epitalon was obtained from Dr G.I. Grigoriev, Institute of Bioregulation and Gerontology, St. Petersburg, Russia.

Animals. Two-month-old outbred male LIO rats from the Animal Department of Central Research Roentgeno-Radiological Institute, St. Petersburg, were used in the study. Animals were kept under a standard light/dark regimen (12:12 h) at $22 \pm 2^\circ\text{C}$. They received standard laboratory chow and tap water *ad libitum*.

Tumorigenic experiments. Eighty rats were exposed weekly to five subcutaneous injections of DMH at a dose of 21 mg/kg body weight (calculated as a base). Under this regimen, the carcinogen induced colon tumors in most of the rats (1). DMH was *ex tempore* dissolved in normal saline and neutralized with sodium bicarbonate (pH 7.0). Rats were divided into four groups. Control rats (group 1) received saline at a dose of 0.1 ml during the entire experiment, lasting 6 months. Rats from groups 2, 3 and 4 were treated subcutaneously with Epitalon at a single dose of 1 μg five times a week. Rats in group 2 were treated with Epitalon for 6 months, from the first injection of DMH till the end of experiment. Rats in group 3 were treated with Epitalon for about 5 months, after termination of carcinogen injections till the end of the experiment. Rats in group 4 were treated with Epitalon only during the period of DMH exposure (for the first 5 weeks of the experiment). All animals were weighed weekly. The experiment was terminated 6 months after the first injection of carcinogen, and all rats were sacrificed by ether vapor.

Pathological investigation. All animals were autopsied. The intestine was opened longitudinally. The position and size of

each tumor were recorded. All tumors and colon mucosa were fixed in 10% neutral formalin and, after routine histological processing, were embedded in paraffin. Sections (3 μm -thick) through the middle part of each tumor were stained with hematoxylin and eosin (H&E). Proliferative activity of the epithelial cells and condition of the stroma were studied in the mucosa adjacent to a tumor (<1 cm away) and located far from the tumor (≥ 3 cm) as well as in the tumors themselves. The distant mucosa served as an internal control. Proliferative activity was evaluated from the whole slide at a magnification of x150. Mitotic index (MI) was calculated as the percentage of mitotic cells per 10,000 μm^2 of mucosa or tumor. The cells were analyzed with an ocular grid, at a magnification of x600. In each sample 1,000-3,000 cells were evaluated. The stromal areas were evaluated using a 10,000 μm^2 micrometric net in 10 randomly chosen fields of tumors and colon mucosa. The rate of lymphoid infiltration was calculated by determine the number of lymphocytes present as a percentage of the total number of cells in the same areas. The relationship between tumorigenesis and apoptosis was studied by comparing the rate of apoptosis in tumors obtained from different rat groups. The apoptotic index (AI) was determined using the ApopTag marker (Oncor, Inc., Gaithersburg, MD) and was calculated as the percentage of terminal deoxynucleotidyl transferase-mediated dUTP-biotin nick end labeling (TUNEL)-positive cells per 1,000 cells (11). All data were analyzed statistically by Student's t-test.

Results

DMH stimulated mitotic activity of the epithelial cells in the mucosa adjacent to a tumor (Fig. 1). In tumors, this parameter was sevenfold that in the mucosal epithelial cells located far from the tumor (Table I). Epitalon inhibited mitotic activity of the epithelial cells adjacent to a tumor and especially that of tumor cells when the treatment was given throughout the experiment (group 2, Table I).

Significant differences were found in the AI of tumors obtained from rats treated under the different regimes: long-term treatment with Epitalon (throughout the experiment, group 2) resulted in a significant increase in the number of apoptotic cells in tumors (Fig. 2; Table II).

Table I. Effects of Epitalon on the rate of mitotic activity in the colon (the number of mitotic cells per 10,000 μm^2 of the stroma, mean \pm SE).

Rat groups	Areas of the mucosa located far from a tumor (A)	Areas of the mucosa adjacent to a tumor (B)	Areas of colon adenocarcinomas (C)
1	0.23 \pm 0.08	0.43 \pm 0.08	1.62 \pm 0.13 ^a
2	0.23 \pm 0.09	0.23 \pm 0.07	1.03 \pm 0.13 ^{a,b}
3	0.25 \pm 0.09	0.22 \pm 0.07	1.36 \pm 0.13 ^a
4	0.22 \pm 0.07	0.40 \pm 0.08	1.66 \pm 0.14 ^{a,c}

All rats were exposed to DMH. Rat groups: 1, Control, non-treated. 2, Treated with Epitalon from the first injection of DMH till the end of experiment, for 6 months. 3, Treated with Epitalon after the last injection of DMH till the end of experiment. 4, Treated with Epitalon only during the exposure to DMH, for the first 5 weeks of experiment. ^aSignificantly different from areas A and B, $p < 0.01$. ^bSignificantly different from group 1, $p < 0.01$. ^cSignificantly different from group 2, $p < 0.01$.




Figure 2. Colon tumors obtained from a control rat (A) and a rat treated with Epitalon (B). Note the abundance of apoptotic cells (arrows) in tumors from the experimental rat. TUNEL reaction. $\times 400$.

Table II. Apoptotic index (AI) in colon tumors obtained from different rat groups (mean \pm SE).

Rat groups	AI
1 ^a	1.64 \pm 0.27
2	12.74 \pm 1.19 ^b
3	2.76 \pm 0.32
4	2.17 \pm 0.23

^aSee footnotes at Table I. ^bSignificantly different from the other groups, $p < 0.01$.

The morphometric study showed that DMH-induced colon carcinogenesis is accompanied by a decrease in the size of the stromal area and in that of the lymphoid infiltration into tumors and in the mucosa adjacent to the tumors, relative to mucosal areas located far from the tumors (Table III). Epitalon attenuated this effect, especially when the treatment

was continued for the entire experimental period. A comparison between the different experimental groups showed that Epitalon increases the stromal areas and the intensity of lymphoid infiltration both in the colon mucosa adjacent to a tumor and in the tumors themselves (Tables III and IV).

Discussion

As previously shown, Epitalon inhibits chemically-induced colon carcinogenesis in rats (5). We found that Epitalon inhibited mitotic activity of the epithelial cells adjacent to a tumor and especially that of the tumor cells themselves when the treatment was given throughout the experiment. Mucosal samples near tumors showed a small, non-significant increase in mitotic activity relative to the more distant samples (12). The mean number of mitoses/crypt has been found to be similar for controls and adenomas but greater for cancers, and particularly for familial adenomatous polyposis (12). Although hyperproliferation of the normal colorectal mucosa in patients with sporadic adenomas has been described (13), no differences were found in the crypt-cell proliferation state between patients with sporadic adenomas and controls (12). Epithelial cell proliferation reflects the progression of carcino-

Table III. Effects of Epitalon on the stromal areas and lymphoid infiltration in the colon (% of the entire area of the mucosal layer at the slide, mean \pm SE).

Rat groups	Areas of the mucosa located far from a tumor (A)		Areas of the mucosa adjacent to a tumor (B)		Areas of colon adenocarcinomas (C)	
	Stroma	Lymph infiltration	Stroma	Lymph infiltration	Stroma	Lymph infiltration
1	26.1 \pm 1.3	14.9 \pm 1.3	21.3 \pm 1.2 ^a	11.5 \pm 0.7	17.0 \pm 1.7 ^a	6.5 \pm 0.9 ^b
2	32.2 \pm 1.0 ^c	16.2 \pm 0.9 ^d	28.0 \pm 2.1 ^c	14.9 \pm 1.6 ^{c,c}	18.3 \pm 1.4 ^b	6.4 \pm 0.8 ^b
3	28.8 \pm 1.3	12.5 \pm 0.9	29.2 \pm 1.4 ^c	11.7 \pm 0.9	17.8 \pm 1.2 ^b	5.4 \pm 0.7 ^b
4	32.2 \pm 1.7 ^c	13.8 \pm 0.4	24.2 \pm 1.2 ^a	9.75 \pm 0.6 ^f	18.7 \pm 1.3 ^b	6.8 \pm 0.7 ^b

Rat groups: see footnotes at Table I. ^aSignificantly different from A, $p < 0.05-0.01$. ^bSignificantly different from A and B, $p < 0.01$. ^cSignificantly different from group 1, $p < 0.05-0.01$. ^dSignificantly different from groups 3 and 4, $p < 0.05$. ^eSignificantly different from groups 1, 3 and 4, $p < 0.05$. ^fSignificantly different from areas A and C, $p < 0.01$.

Table IV. Effects of Epitalon on the intensity of lymphoid infiltration in the colon (the number of lymphoid cells per 10,000 μm^2 of the stroma, mean \pm SE).

Rat groups	Areas of the mucosa located far from a tumor (A)	Areas of the mucosa adjacent to a tumor (B)	Areas of colon adenocarcinomas (C)
1	132.3 \pm 10.7	100.5 \pm 3.9 ^a	41.1 \pm 4.9 ^b
2	135.2 \pm 9.8	121.7 \pm 10.9	53.3 \pm 6.8 ^b
3	95.9 \pm 9.2 ^c	97.3 \pm 4.6	37.3 \pm 4.8 ^b
4	101.3 \pm 4.0 ^c	101.0 \pm 5.1	42.1 \pm 6.0 ^b

Rat groups: see footnotes at Table I. ^aSignificantly different from A, $p < 0.05$. ^bSignificantly different from A and B, $p < 0.01$. ^cSignificantly different from groups 1 and 2, $p < 0.05$.

genesis. The histogenesis of hyperplastic polyps with atypia involves the hyperplastic polyp-carcinoma sequence, whereas the development of tubulovillous or serrated adenomas may involve the tubulovillous adenoma-carcinoma or serrated adenoma-carcinoma sequence (14).

Long-term treatment with Epitalon resulted in a significant increase in the number of apoptotic cells in tumors. Similar data were obtained in our previous study (9). We observed an increase in the AI of colon tumors in rats exposed to DMH alone as compared with the AI in the colon mucosa of the same rats, as well as controls (11). It has been suggested that apoptosis is inhibited during the progression of colon tumorigenesis (15). In azoxymethane-induced rat colon carcinogenesis, a decrease in the expression of the apoptotic repressor bcl-2 and an increase in the expression of the apoptosis accelerator, bax protein, have been shown (16). Our data appear to be relevant to this observation.

Cell kinetic status affects the macroscopic morphology of colorectal neoplasms (17). AI is closely associated with macroscopic morphology in adenomas but not in carcinomas. The MI is relatively constant among the three macroscopic types in adenomas and carcinomas, although overall MI is

significantly higher in carcinomas than in adenomas, whereas AI does not differ. The macroscopic morphology of colorectal adenomas is determined by the apoptosis and not by the proliferation, and high apoptosis found in depressed adenomas implies low net growth.

We have observed a significant decrease in the mean square of stromal areas and of lymphoid infiltrates in the colon mucosa adjacent to a tumor and in tumors of rats exposed to DMH alone, as compared to intact controls. The treatment with Epitalon was followed by a reduction in these parameters. The strongest inhibitory effect of Epitalon occurred when the treatment was continued throughout the experimental period (group 2).

Similar data have been obtained in our previous study on the effect of melatonin on proliferative activity in the colon mucosa and colon tumors (9). The multiplicity of DMH-induced colon tumors in rats was significantly reduced after treatment with melatonin and this effect was correlated with an inhibition of mitosis and stimulation of apoptosis. These data are in agreement with observations of the inhibitory effect of melatonin on proliferative activity in rodent colon cells (7), grafted hepatoma cells (18) and human peripheral

blood lymphocytes (19). Melatonin decreased cell proliferation, and increased the AI in colon cancer cells (2). Moreover, melatonin significantly decreased the proliferation-to-apoptosis ratio, an additional parameter confirming the inhibition of tumor growth (2).

The finding in the present study is in accordance with our previous observations that the inhibitory effect of Epitalon on colon cancer is manifested at various stages of carcinogenesis (5,6) and especially when it is applied during the entire process of tumor development. It is suggested that Epitalon can act both directly and indirectly through the stimulation of pineal function (20). Our data confirm the hypothesis of the preventive role of the pineal gland in cancer development, realized through the action of melatonin as well as of the pineal peptides.

Acknowledgments

This work was supported in part by grant #00-04.48481 from The Russian Foundation for Basic Research (Russia) and by grant #039-6119 from the Israeli Government.

References

- Anisimov VN, Popovich IG and Zabezhinski MA: Melatonin and colon carcinogenesis: I. Inhibitory effect of melatonin on development of intestinal tumors induced by 1,2-dimethylhydrazine in rats. *Carcinogenesis* 18: 1549-1553, 1997.
- Winczyk K, Pawlikowski M, Lawnicka H, Kunert-Radek J, Spadoni G, Tarzia G and Karasek M: Effects of melatonin and melatonin receptors ligand N-[(4-methoxy-1H-indol-2-yl)methyl]propanamide on murine colon 38 cancer growth *in vitro* and *in vivo*. *Neuroendocrinol Lett* 23: 50-54, 2002.
- Khavinson VKh, Morozov VG and Anisimov VN: Experimental studies of the pineal gland preparation Epithalamin. In: *The Pineal Gland and Cancer*. Bartsch C, Bartsch H, Blask DE, Cardinali DP, Hrushesky WJM and Mecke D (eds). Springer Verlag, Berlin, pp294-306, 2001.
- Anisimov VN: Mutant and genetically modified mice as models for studying the relationship between aging and carcinogenesis. *Mech Aging Dev* 122: 1221-1255, 2001.
- Anisimov VN, Khavinson VKh, Popovich IG and Zabezhinski MA: Inhibitory effect of peptide Epitalon on colon carcinogenesis induced by 1,2-dimethylhydrazine in rats. *Cancer Lett* 183: 1-8, 2002.
- Zusman I, Reifen R, Livni O, Smimoff P, Gurevich P, Sandler B, Nyska A, Gal R, Tendler Y and Madar Z: Role of apoptosis, proliferating cell nuclear antigen and p53 protein in chemically-induced colon cancer in rats fed corn cob fiber treated with the fungus *Pleurotus ostreatus*. *Anticancer Res* 17: 2105-2114, 1997.
- Lewinski A, Rybicka I, Wajs E, Szkudlinski M and Pawlikowski M: Influence of pineal indolamines on the mitotic activity of gastric and colonic mucosa epithelial cells in the rat: interaction with omeprazole. *J Pineal Res* 10: 104-108, 1991.
- Anisimov VN, Popovich IG, Shtylik AV, Zabezhinski MA, Ben-Hur H, Gurevich P, Berman V, Tendler Y and Zusman I: Melatonin and colon carcinogenesis. III. Effect of melatonin on proliferative activity and apoptosis in colon mucosa and colon tumors induced by 1,2-dimethylhydrazine in rats. *Toxicol Exp Pathol* 52: 71-76, 2000.
- Kossov G, Ben-Hur H, Popovich IG, Zabezhinski M, Anisimov V and Zusman I: Melatonin and colon carcinogenesis. IV. Effect of melatonin on proliferative activity and expression of apoptosis-related proteins in the spleen of the rats exposed to 1,2-dimethylhydrazine. *Oncol Rep* 7: 1401-1405, 2000.
- Anisimov VN: Melatonin and colon carcinogenesis. In: *The Pineal Gland and Cancer*. Bartsch C, Bartsch H, Blask DE, Cardinali DP, Hrushesky WJM and Mecke D (eds). Springer, Berlin, Heidelberg, pp240-258, 2001.
- Tendler Y, Ben-Hur H, Gurevich P, Zion H, Berman V, Sandler B, Nyska A, Zinder O and Zusman I: Role of apoptosis, proliferating cell nuclear antigen and p53 protein in the immune response of the rat colon cells to cancer and vaccination with the anti-p53 polyclonal antibodies. *Anticancer Res* 17: 4653-4658, 1997.
- Mills SJ, Mathers JC, Chapman PD, Burn J and Gunn A: Colonic crypt cell proliferation state assessed by whole crypt microdissection in sporadic neoplasia and familial adenomatous polyposis. *Gut* 48: 41-46, 2001.
- Anti M, Marra G and Armelao F: Rectal epithelial cell proliferation patterns as predictors of adenomatous colorectal polyp recurrence. *Gut* 34: 525-530, 1993.
- Hamatani S, Wada R, Tsujimoto S, Yanagita M, Mitsuda A, Hasegawa C, Kuwahata N, Yokouchi Y and Miura M: Histomorphometric characteristics and cellular kinetics of colorectal polyps with epithelial serrated proliferation adjacent to carcinoma. *Oncol Rep* 8: 49-55, 2001.
- Bedi A, Pasricha PJ, Akhtar AJ, Barber JP, Bedi GC, Giardiello FM, Zehnbauer BA, Hamilton SR and Jones RJ: Inhibition of apoptosis during development of colorectal cancer. *Cancer Res* 55: 1811-1816, 1995.
- Hirose Y, Yoshimi N, Suzui M, Kawabata K, Tanaka T and Mori H: Expression of bcl-2, bax, and bcl-X_L proteins in azoxymethane-induced rat colonic adenocarcinomas. *Mol Carcinog* 19: 25-30, 1997.
- Nomura M, Watari J, Yokota K, Saitoh Y, Obara T and Kohgo Y: Morphogenesis of non-polypoid colorectal adenomas and early carcinomas assessed by proliferation cell and apoptosis. *Virchows Arch* 437: 17-24, 2000.
- Karasek M, Liberski P, Kunert-Radek L and Bartkowiak J: Influence of melatonin on the proliferation of hepatoma cells in the Syrian hamster: *in vivo* and *in vitro* study. *J Pineal Res* 13: 107-110, 1992.
- Vijayalax MI, Reiter RJ, Leal BZ and Meltz ML: Effect of melatonin on mitotic and proliferation indices, and sister chromatid exchange in human blood lymphocytes. *Mutat Res* 351: 187-192, 1996.
- Khavinson VKh, Yakovleva ND, Popuchiev VV, Kvetnoy IM and Manokhina RP: Reparative action of Epitalon on the ultrastructure of pineal gland in irradiated rats. *Bull Exp Biol Med* 131: 98-103, 2001.